

ANOMALOUS COUPLINGS OF NON-BPS D-BRANES

Marco Billó

*Dipartimento di Fisica Teorica, Università di Torino and
I.N.F.N., Sezione di Torino, via P. Giuria 1, I-10125, Torino, Italy*

Ben Craps and Frederik Roose

*Instituut voor Theoretische Fysica
Katholieke Universiteit Leuven, B-3001 Leuven, Belgium*

Abstract Non-BPS type II D-branes couple to R-R potentials via an action that, upon tachyon condensation, gives rise to the Wess-Zumino action of BPS D-branes.

Non-BPS branes in string theory have received a lot of attention after the seminal work by Sen [1], as such non-BPS states allow for highly non-trivial tests of string dualities. In a related development, unstable and/or non-BPS D-branes have played a key role in the unifying description of all lower-dimensional D-branes as non-trivial excitations on suitable configurations of higher-dimensional branes. This description exhibits in a constructive way the fact that D-brane charges take values in appropriate K-theory groups of space-time.

For type IIB this was demonstrated by Witten in [2], where all branes were built from sufficiently many D9–anti-D9 pairs. In this set-up, the D p -brane that emerges via a process of tachyon condensation inherits its anomalous Wess-Zumino couplings to the R-R fields [3],

$$S_{\text{WZ}} = \frac{T_p}{\kappa} \int_{p+1} C \wedge \text{Tr} e^{2\pi\alpha' F+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)}, \quad (1.1)$$

from the analogous couplings of the parent branes. In eq. (1.1), T_p/κ denotes the D p -brane tension, C a formal sum of R-R potentials, F the gauge field on the brane and B the NS-NS two-form. The trace is over the Chan-Paton indices. Further, R_T and R_N are the curvatures

of the tangent and normal bundles of the D-brane world-volume, and $\hat{A}$ denotes the A-roof genus.

As for type IIA, Horava described in [4] how to construct BPS $D(p - 2k - 1)$ -branes as bound states of (sufficiently many) unstable Dp -branes (and thus how to construct all type IIA D-branes in terms of D9-branes). The lower-dimensional BPS branes arise as the result of the condensation of a tachyon field into a vortex configuration, accompanied by non-trivial gauge fields. In this case, it was not clear how the lower-dimensional BPS branes acquire the R-R couplings of eq. (1.1).

In [5] we argued that all type II non-BPS branes couple universally to Ramond-Ramond fields as given by¹

$$S'_{\text{WZ}} = a \int_{p+1} C \wedge d \text{Tr } T e^{2\pi\alpha' F+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)} , \quad (1.2)$$

where T is the real, adjoint tachyon field living on the non-BPS brane and a is a constant. Let us indicate how the R-R couplings of eq. (1.2) account for the R-R couplings (1.1) of the stable lower-dimensional brane that emerges from the process of tachyon condensation. The cases $Dp \rightarrow D(p - 1)$ and $Dp \rightarrow D(p - 3)$ will be treated in detail. It will turn out, for instance, that the R-R charges of the D8-branes and D6-branes one constructs from unstable D9-branes [4] have the expected ratio.

Consider first a single non-BPS Dp -brane. There is a real tachyon field living on its worldvolume. The tachyon potential is assumed to be such that the vacuum manifold consists of the two points $\{T_0, -T_0\}$. The tachyon can condense in a non-trivial (anti)-kink configuration $T(x)$ depending on a single coordinate. The R-R coupling (1.2) on the Dp -brane reads in this case

$$a \int_{p+1} C \wedge dT \wedge e^{2\pi\alpha' F+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)} . \quad (1.3)$$

It involves the topological density $\partial_x T(x)$, which is localized at the core of the kink and is such that $\int dT(x) = \pm 2T_0$. In the limit of zero size, $dT(x) = 2T_0 \delta(x - x_0) dx$, and the above action takes the form of the usual Wess-Zumino effective action for a BPS $D(p - 1)$ -brane, localized in the x -direction at x_0 :

$$2T_0 a \int_p C \wedge e^{2\pi\alpha' F+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)} . \quad (1.4)$$

¹One term of this action (the one describing the coupling of a non-BPS D9-brane to C_9) was conjectured to be present in Ref. [4] and obtained by a disc computation in Ref. [1].

As a less trivial example, let us start from two unstable Dp-branes. The tachyon field T , transforming in the adjoint of the U(2) gauge group, can form a non-trivial vortex configuration in co-dimension three. The tachyon potential is assumed to be such that the minima of T have the eigenvalues $(T_0, -T_0)$, so that the vacuum manifold is $\mathcal{V} = \text{U}(2)/(\text{U}(1) \times \text{U}(1)) = S^2$. The possible stable vortex configurations $T(\mathbf{x})$, depending on 3 coordinates x^i transverse to the $(p-2)$ -dimensional core of the vortex, are classified by the non-trivial embeddings of the “sphere at infinity” S_∞^2 into the vacuum manifold, namely by $\pi_2(\mathcal{V}) = \mathbf{Z}$.

Apart from the “center of mass” U(1) subgroup, we are in the situation of the Georgi-Glashow model, where the tachyon field $T(\mathbf{x}) = T^a(\mathbf{x})\sigma^a$ (σ^a being the Pauli matrices) sits in the adjoint of SU(2), and the vacuum manifold is described by $T^a T^a = T_0^2$. The vortex configuration of winding number one of the tachyon, which is the ’t Hooft-Polyakov monopole, is accompanied by a non-trivial SU(2) gauge field:

$$\begin{aligned} T(\mathbf{x}) &= f(r)\sigma_a x^a, \\ \mathcal{A}_i^a(\mathbf{x}) &= h(r)\epsilon_{ij}^a x^j, \end{aligned} \quad (1.5)$$

where r is the radial distance in the three transverse directions; the prefactors $f(r)$ and $h(r)$ go to constants for $r \rightarrow 0$, while $f(r) \sim T_0/r$ and $h(r) \sim 1/r^2$ for $r \rightarrow \infty$. The field-strength in the unbroken U(1) direction,

$$\mathcal{G}_{ij} = \frac{T^a}{T_0} \mathcal{F}_{ij}^a, \quad (1.6)$$

corresponds to a non-trivial U(1) bundle on the sphere at infinity, *i.e.* the magnetic charge $g = \int_{S_\infty^2} \mathcal{G}$ is non-zero (and in fact equals the winding number of the vortex in appropriate units). The magnetic charge density in the transverse directions, defined by $d\mathcal{G} = \rho(\mathbf{x})d^3x$, is concentrated at the core of the vortex solution. In the zero size limit, there is a point-like magnetic charge at the location of the core: $\rho(\mathbf{x}) = g\delta^3(\mathbf{x} - \mathbf{x}^0)$.

The WZ action (1.2) for the Dp-brane can be rewritten as

$$a \int_{p+1} C \wedge d\text{Tr}\{T e^{2\pi\alpha'\mathcal{F}}\} \wedge e^{2\pi\alpha'\hat{F}+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)}, \quad (1.7)$$

where we have split the U(2) field-strength into its SU(2) part $\mathcal{F}$ and its U(1) part $\hat{F}$. Inserting the ’t Hooft-Polyakov configuration for the tachyon and the SU(2) gauge field, we see that Eq. (1.7) involves the curl $d\mathcal{G}$ of the magnetic monopole field $\mathcal{G} = T^a \mathcal{F}^a$; we end up with

$$2\pi\alpha' a T_0 \int_{p+1} C \wedge \rho(\mathbf{x})d^3x \wedge e^{2\pi\alpha'\hat{F}+B} \wedge \sqrt{\hat{A}(R_T)/\hat{A}(R_N)}. \quad (1.8)$$

We have a distribution of $D(p-3)$ -brane charge localized at the core of the vortex; in particular, in the limit of zero-size core we recover the R-R couplings (1.2) of a BPS $D(p-3)$ -brane that supports the $U(1)$ gauge field $\hat{F}$.

Since the minimal magnetic charge g is 4π in our units, Eqs. (1.4) and (1.8) lead to the expected ratio $4\pi^2\alpha'$ for the R-R charges of $D(p-3)$ - and $D(p-1)$ -branes.

The mechanism described above generalizes to the reduction of a non-BPS Dp -brane to a $D(p-2k-1)$ -brane via tachyon condensation, described in [4]. In this case, it is convenient to start with 2^k unstable Dp -branes. The tachyon field, which sits in the adjoint of $U(2^k)$, can be in a vortex configuration, accompanied by non-trivial gauge fields, such that there is a non-zero generalized magnetic charge $\int_{S_\infty^{2k}} \text{Tr}\{T(\mathcal{F})^k\}$. The WZ action (1.2) then contains the generalized magnetic charge density $d\text{Tr}\{T\mathcal{F}^k\} = \rho(\mathbf{x})d^{2k+1}x$, localized at the core of the vortex. In the zero-size limit, we are left with the WZ action for a $D(p-2k-1)$ -brane.

The form of the R-R couplings (1.2) can be checked directly by computing the corresponding string scattering amplitudes [5].

Acknowledgments

B.C. and F.R. would like to thank the organisers for a very nice school, and for financial support. This work was supported by the European Commission TMR programme ERBFMRX-CT96-0045. B.C. is Aspirant FWO-Vlaanderen.

References

- [1] Sen, A. (1998) *JHEP* **9809** 023, [hep-th/9808141](#).
- [2] Witten, E. (1998) D-branes and K-theory, *JHEP* **9812** 019, [hep-th/9810188](#).
- [3] Green, M., Harvey, J.A. and Moore, G. (1997) I-brane inflow and anomalous couplings on D-branes, *Class.Quant.Grav.* **14**, pp. 47-52, [hep-th/9605033](#);
Cheung, Y.K. and Yin, Z. (1998) Anomalies, branes, and currents, *Nucl. Phys.* **B517**, pp. 69-91, [hep-th/9710206](#).
- [4] Horava, P. (1998) Type IIA D-branes, K theory, and Matrix theory, *Adv. Theor. Math. Phys.* **2** pp. 1373-1404, [hep-th/9812135](#).
- [5] Billó, M., Craps, B. and Roose, F. (1999) Ramond-Ramond couplings of non-BPS D-branes, *JHEP* **9906** 033, [hep-th/9905157](#).